

# The Ehrhart polynomial of the Birkhoff polytope <sup>1</sup>

MATTHIAS BECK AND DENNIS PIXTON

*All means (even continuous) sanctify the discrete end.* Doron Zeilberger<sup>2</sup>

*Abstract:* The  $n^{\text{th}}$  Birkhoff polytope is the set of all doubly stochastic  $n \times n$  matrices, that is, those matrices with nonnegative real coefficients in which every row and column sums to one. A wide open problem concerns the volumes of these polytopes, which have been known for  $n \leq 8$ . We present a new, complex-analytic way to compute the Ehrhart polynomial of the Birkhoff polytope, that is, the function counting the integer points in the dilated polytope. One reason to be interested in this counting function is that the leading term of the Ehrhart polynomial is—up to a trivial factor—the volume of the polytope. We implemented our methods in the form of a computer program, which yielded the Ehrhart polynomial (and hence the volume) of the ninth Birkhoff polytope.

## 1 Introduction

One of the most intriguing objects of combinatorial geometry is the  $n^{\text{th}}$  Birkhoff polytope

$$\mathcal{B}_n = \left\{ \left( \begin{array}{ccc} x_{11} & \cdots & x_{1n} \\ \vdots & & \vdots \\ x_{n1} & \cdots & x_{nn} \end{array} \right) \in \mathbb{R}^{n^2} : x_{jk} \geq 0, \sum_j x_{jk} = 1 \text{ for all } 1 \leq k \leq n \right. \\ \left. \sum_k x_{jk} = 1 \text{ for all } 1 \leq j \leq n \right\},$$

often described as the set of all  $n \times n$  doubly stochastic matrices.  $\mathcal{B}_n$  is a convex polytope with integer vertices. It possesses fascinating combinatorial properties [4, 5, 6, 8, 23] and relates to many mathematical areas [10, 14]. A long-standing open problem is the determination of the relative volume of  $\mathcal{B}_n$ , which had been known only up to  $n = 8$  [7, 17]. In this paper, we propose a new method of calculating this volume and use it to compute  $\text{vol} \mathcal{B}_9$ .

One of the recent attempts to compute  $\text{vol} \mathcal{B}_n$  relies on the theory of counting functions for the integer points in polytopes. Ehrhart proved [11] that for a polytope  $\mathcal{P} \subset \mathbb{R}^d$  with integral vertices, the number

$$L_{\mathcal{P}}(t) := \#(t\mathcal{P} \cap \mathbb{Z}^d)$$

is a polynomial in the positive integer variable  $t$ . He showed various other properties for this counting function (in fact, in the more general setting of  $\mathcal{P}$  having rational vertices), of which we mention three here:

- The degree of  $L_{\mathcal{P}}$  is the dimension of  $\mathcal{P}$ .
- The leading term of  $L_{\mathcal{P}}$  is the relative volume of  $\mathcal{P}$ , normalized with respect to the sublattice of  $\mathbb{Z}^d$  on the affine subspace spanned by  $\mathcal{P}$ .

---

<sup>1</sup>To appear in *Discrete & Computational Geometry*

<sup>2</sup>E-mail to the first author on Wed, 14 Aug 2002 16:29:35 -0400 (EDT)

- Since  $L_{\mathcal{P}}$  is a polynomial, we can evaluate it at nonpositive integers. These evaluations yield

$$L_{\mathcal{P}}(0) = \chi(\mathcal{P}) , \tag{1}$$

$$L_{\mathcal{P}}(-t) = (-1)^{\dim \mathcal{P}} L_{\mathcal{P}^\circ}(t) . \tag{2}$$

Here  $\chi(\mathcal{P})$  denotes the Euler characteristic,  $\mathcal{P}^\circ$  the relative interior of  $\mathcal{P}$ . The reciprocity law (2) was in its full generality proved by Macdonald [15].

The application of this theory to the Birkhoff polytope  $\mathcal{B}_n$  incorporates the nice interpretation of the number of integral points in  $t\mathcal{B}_n$  as the number of *semi-magic squares*, namely, square matrices whose nonnegative integral coefficients sum up to the same integer  $t$  along each row and column.

We will denote the Ehrhart polynomial of  $\mathcal{B}_n$  by

$$H_n(t) := L_{\mathcal{B}_n}(t) .$$

It is not hard to see that  $\dim \mathcal{B}_n = (n-1)^2$ , hence  $H_n$  is a polynomial in  $t$  of degree  $(n-1)^2$ . The first two of these polynomials are trivial:

$$H_1(t) = 1 , \quad H_2(t) = t + 1 ,$$

the first nontrivial case was computed by MacMahon [16] as

$$H_3(t) = 3 \binom{t+3}{4} + \binom{t+2}{2} . \tag{3}$$

The structural properties of  $H_n$  were first studied in [12, 18, 20]. It is a nice exercise to deduce from (2) that

$$H_n(-n-t) = (-1)^{n-1} H_n(t) \tag{4}$$

and

$$H_n(-1) = H_n(-2) = \dots = H_n(-n+1) = 0 .$$

This allows the following strategy of computing  $H_n$ , and therefore, the volume of  $\mathcal{B}_n$ : compute the first  $\binom{n-1}{2}$  values of  $H_n$ , use the above symmetry and trivial values of  $H_n$ , and calculate the polynomial  $H_n$  by interpolation. In fact, as far as we are aware of, the volume of  $\mathcal{B}_8$  was computed using essentially this method, combined with some nice computational tricks [7, 17].

We propose a new, completely different approach of computing  $H_n$  (and hence  $\text{vol } \mathcal{B}_n$ ). It is based on an analytic method by the first author of computing the Ehrhart polynomial of a polytope [2]. We will introduce the application of this method to the Birkhoff polytope in the following section.

Some recent refreshing approaches—of a more algebraic-geometric/topological flavor—to the problem of computing  $\text{vol } \mathcal{B}_n$  and  $H_n$  can be found in [1, 9, 22].



is a  $(2n \times n^2)$ -matrix and  $\mathbf{b} = (1, \dots, 1) \in \mathbb{Z}^{2n}$ . Hence Theorem 1 gives for this special case

$$H_n(t) = \frac{1}{(2\pi i)^{2n}} \int \cdots \int \frac{(z_1 \cdots z_{2n})^{-t-1}}{(1 - z_1 z_{n+1})(1 - z_1 z_{n+2}) \cdots (1 - z_n z_{2n})} dz .$$

Here it is understood that each integral is over a circle with radius  $< 1$  centered at 0; all appearing radii should be different. We can separate, say, the last  $n$  variables and obtain

$$H_n(t) = \frac{1}{(2\pi i)^n} \int \cdots \int (z_1 \cdots z_n)^{-t-1} \left( \frac{1}{2\pi i} \int \frac{z^{-t-1}}{(1 - z_1 z) \cdots (1 - z_n z)} dz \right)^n dz_n \cdots dz_1 .$$

We may choose the radius of the integration circle of the innermost integral to be smaller than the radii of the other integration paths. Then this innermost integral is easy to compute: It is equal to the residue at 0 of

$$\frac{1}{z^{t+1}(1 - z_1 z) \cdots (1 - z_n z)}$$

and, by the residue theorem, equal to the negative of the sum of the residues at  $z_1^{-1}, \dots, z_n^{-1}$ . (Note that here we use the fact that  $t > 0$ .) The residues at these simple poles are easily computed: the one, say, at  $1/z_1$  can be calculated as

$$\begin{aligned} \lim_{z \rightarrow 1/z_1} \left( z - \frac{1}{z_1} \right) \frac{1}{z^{t+1}(1 - z_1 z) \cdots (1 - z_n z)} &= \lim_{z \rightarrow 1/z_1} \frac{z - \frac{1}{z_1}}{1 - z_1 z} \frac{1}{z^{t+1}(1 - z_2 z) \cdots (1 - z_n z)} \\ &= -\frac{1}{z_1} \frac{z_1^{t+1}}{(1 - z_2/z_1) \cdots (1 - z_n/z_1)} = -\frac{z_1^{t+n-1}}{(z_1 - z_2) \cdots (z_1 - z_n)} . \end{aligned}$$

This yields the starting point for our computations.

**Theorem 2** For any distinct  $0 < \epsilon_1, \dots, \epsilon_n < 1$ ,

$$H_n(t) = \frac{1}{(2\pi i)^n} \int_{|z_1|=\epsilon_1} \cdots \int_{|z_n|=\epsilon_n} (z_1 \cdots z_n)^{-t-1} \left( \sum_{k=1}^n \frac{z_k^{t+n-1}}{\prod_{j \neq k} (z_k - z_j)} \right)^n dz_n \cdots dz_1 .$$

*Remark.* It can be proved from the form of the integrand that  $H_n$  is indeed a polynomial in  $t$ : To compute the integral, one has to execute a (huge) number of limit computations, which yield “at worst” powers of  $t$  (as a consequence of L’Hospital’s Rule). In fact, one can make this property more apparent by noticing that the expression in parenthesis is actually a polynomial; namely

$$H_n(t) = \frac{1}{(2\pi i)^n} \int_{|z_1|=\epsilon_1} \cdots \int_{|z_n|=\epsilon_n} (z_1 \cdots z_n)^{-t-1} \left( \sum_{m_1 + \cdots + m_n = t} z_1^{m_1} \cdots z_n^{m_n} \right)^n dz_n \cdots dz_1 ,$$

where the sum is over all ordered partitions of  $t$ . This formula can also be proved “more directly” combinatorially.<sup>3</sup>

<sup>3</sup>The authors thank Sinai Robins and Frank Sottile for their help in the proof of this equivalence and its combinatorial interpretation.

### 3 Small $n$ do not require a computer

We will now illustrate the computation of  $H_n$  (and hence  $\text{vol } \mathcal{B}_n$ ) by means of Theorem 2 for  $n = 3$  and 4. These calculations “by hand” give an idea what computational tricks one might use in tackling larger  $n$  with the aid of a computer.

By the theorem,

$$H_3(t) = \frac{1}{(2\pi i)^3} \int (z_1 z_2 z_3)^{-t-1} \left( \frac{z_1^{t+2}}{(z_1 - z_2)(z_1 - z_3)} + \frac{z_2^{t+2}}{(z_2 - z_1)(z_2 - z_3)} + \frac{z_3^{t+2}}{(z_3 - z_1)(z_3 - z_2)} \right)^3 dz.$$

We have to order the radii of the integration paths for each variable; we choose  $0 < \epsilon_3 < \epsilon_2 < \epsilon_1 < 1$ . We heavily use this fact after multiplying out the cubic: integrating, for example, the term

$$\frac{z_1^{-t-1} z_2^{-t-1} z_3^{2t+5}}{(z_3 - z_2)^3 (z_3 - z_1)^3}$$

with respect to  $z_3$  gives 0, as this function is analytic at the  $z_3$ -origin and  $|z_1|, |z_2| > \epsilon_3$ . After exploiting this observation for all the terms stemming from the cubic, the only integrals surviving are

$$\frac{1}{(2\pi i)^3} \int \frac{z_1^{2t+5} z_2^{-t-1} z_3^{-t-1}}{(z_1 - z_2)^3 (z_1 - z_3)^3} dz$$

and

$$- \frac{3}{(2\pi i)^3} \int \frac{z_1^{t+3} z_2 z_3^{-t-1}}{(z_1 - z_2)^3 (z_1 - z_3)^2 (z_2 - z_3)} dz.$$

The first integral factors and yields, again by residue calculus,

$$\begin{aligned} \frac{1}{(2\pi i)^3} \int \frac{z_1^{2t+5} z_2^{-t-1} z_3^{-t-1}}{(z_1 - z_2)^3 (z_1 - z_3)^3} dz &= \frac{1}{(2\pi i)^3} \int z_1^{2t+5} \left( \int \frac{z^{-t-1}}{(z_1 - z)^3} dz \right)^2 dz_1 \\ &= \frac{1}{2\pi i} \int z_1^{2t+5} \left( -\frac{1}{2}(-t-1)(-t-2) z_1^{-t-3} \right)^2 dz_1 = \binom{t+2}{2}. \end{aligned}$$

For the second integral, it is most efficient to integrate with respect to  $z_2$  first:

$$\begin{aligned} - \frac{3}{(2\pi i)^3} \int \frac{z_1^{t+3} z_2 z_3^{-t-1}}{(z_1 - z_2)^3 (z_1 - z_3)^2 (z_2 - z_3)} dz &= - \frac{3}{(2\pi i)^2} \int \frac{z_1^{t+3} z_3^{-t}}{(z_1 - z_3)^5} dz_3 dz_1 \\ &= - \frac{3}{2\pi i} \int z_1^{t+3} \frac{1}{4!} (-t)(-t-1)(-t-2)(-t-3) z_1^{-t-4} dz_1 = -3 \binom{t+3}{4}. \end{aligned}$$

Adding up the last two lines gives finally

$$H_3(t) = \binom{t+2}{2} - 3 \binom{t+3}{4} = \frac{1}{8}t^4 + \frac{3}{4}t^3 + \frac{15}{8}t^2 + \frac{9}{4}t + 1,$$

which is equal to (3). To obtain the volume of  $\mathcal{B}_3$ , the leading term of  $H_3$  has to be multiplied by the relative volume of the fundamental domain of the sublattice of  $\mathbb{Z}^9$  in the affine space spanned by  $\mathcal{B}_3$ . This volume is 9; hence

$$\text{vol } \mathcal{B}_3 = \frac{9}{8}.$$

In general, it is not hard to prove (see, for example, the appendix of [7]) that the relative volume of the fundamental domain of the sublattice of  $\mathbb{Z}^{n^2}$  in the affine space spanned by  $\mathcal{B}_n$  is  $n^{n-1}$ .

The number of integrals we have to evaluate to compute  $H_4$  is only slightly higher. By Theorem 2,

$$H_4(t) = \frac{1}{(2\pi i)^4} \int_{|z_1|=\epsilon_1} \int_{|z_2|=\epsilon_2} \int_{|z_3|=\epsilon_3} \int_{|z_4|=\epsilon_4} (z_1 z_2 z_3 z_4)^{-t-1} \left( \sum_{k=1}^4 \frac{z_k^{t+3}}{\prod_{j \neq k} (z_k - z_j)} \right)^4 dz.$$

Again we have a choice of ordering the radii; we use  $0 < \epsilon_4 < \epsilon_3 < \epsilon_2 < \epsilon_1 < 1$ . After multiplying out the quartic, we have to calculate five integrals; their evaluation—again straightforward by means of the residue theorem—is as follows. As before, we can ‘save’ computation effort by choosing a particular order with which we integrate and by factoring an integral if possible.

$$\begin{aligned} & \frac{1}{(2\pi i)^4} \int \frac{z_1^{3t+11} z_2^{-t-1} z_3^{-t-1} z_4^{-t-1}}{(z_1 - z_2)^4 (z_1 - z_3)^4 (z_1 - z_4)^4} dz = \frac{1}{(2\pi i)^4} \int z_1^{3t+11} \left( \int \frac{z^{-t-1}}{(z_1 - z)^4} dz \right)^3 dz_1 \\ & = \binom{t+3}{3}^3, \\ & - \frac{4}{(2\pi i)^4} \int \frac{z_1^{2t+8} z_2^2 z_3^{-t-1} z_4^{-t-1}}{(z_1 - z_2)^4 (z_1 - z_3)^3 (z_1 - z_4)^3 (z_2 - z_3)(z_2 - z_4)} dz \\ & = - \frac{4}{(2\pi i)^4} \int \frac{z_1^{2t+8} z_2^2}{(z_1 - z_2)^4} \left( \int \frac{z^{-t-1}}{(z_1 - z)^3 (z_2 - z)} dz \right)^2 dz_1 dz_2 \\ & = \frac{4}{(2\pi i)^2} \int \frac{z_1^{2t+8} z_2^{-t+1}}{(z_1 - z_2)^7} \left( 2 \binom{t+2}{2} \frac{z_1^{-t-3}}{z_1 - z_2} + 2(t+1) \frac{z_1^{-t-2}}{(z_1 - z_2)^2} + 2 \frac{z_1^{-t-1}}{(z_1 - z_2)^3} - \frac{z_2^{-t-1}}{(z_1 - z_2)^3} \right) dz \\ & = 8 \binom{t+2}{2} \binom{t+5}{7} + 8(t+1) \binom{t+6}{8} + 8 \binom{t+7}{9} - 4 \binom{2t+8}{9}, \\ & \frac{4}{(2\pi i)^4} \int \frac{z_1^{2t+8} z_2^{-t-1} z_3^2 z_4^{-t-1}}{(z_1 - z_2)^3 (z_1 - z_3)^4 (z_1 - z_4)^3 (z_2 - z_3)(z_3 - z_4)} dz \\ & = \frac{4}{(2\pi i)^3} \int \frac{z_1^{2t+8} z_2^{-t+1} z_4^{-t-1}}{(z_1 - z_2)^3 (z_1 - z_4)^7 (z_2 - z_4)} dz \\ & = \frac{4}{(2\pi i)^2} \int \frac{z_1^{2t+8} z_4^{-t-1}}{(z_1 - z_4)^7} \left( \binom{t+2}{2} \frac{z_1^{-t-3}}{z_1 - z_4} + (t+1) \frac{z_1^{-t-2}}{(z_1 - z_4)^2} + \frac{z_1^{-t-1}}{(z_1 - z_4)^3} \right) dz \\ & = 4 \left( \binom{t+2}{2} \binom{t+5}{7} + (t+1) \binom{t+6}{8} + \binom{t+7}{9} \right), \\ & \frac{6}{(2\pi i)^4} \int \frac{z_1^{t+5} z_2^{t+5} z_3^{-t-1} z_4^{-t-1}}{(z_1 - z_2)^4 (z_1 - z_3)^2 (z_1 - z_4)^2 (z_2 - z_3)^2 (z_2 - z_4)^2} dz \\ & = \frac{6}{(2\pi i)^4} \int \frac{z_1^{t+5} z_2^{t+5}}{(z_1 - z_2)^4} \left( \int \frac{z^{-t-1}}{(z_1 - z)^2 (z_2 - z)^2} dz \right)^2 dz_1 dz_2 \\ & = \frac{6}{(2\pi i)^2} \int \frac{z_1^{t+5} z_2^{t+5}}{(z_1 - z_2)^4} \left( (t+1)^2 \frac{z_2^{-2t-4}}{(z_1 - z_2)^4} - 4(t+1) \frac{z_2^{-2t-3}}{(z_1 - z_2)^5} + 4 \frac{z_2^{-2t-2}}{(z_1 - z_2)^6} \right) dz \\ & = 6(t+1)^2 \binom{t+5}{7} - 24(t+1) \binom{t+5}{8} + 24 \binom{t+5}{9}, \end{aligned}$$

$$\begin{aligned}
& -\frac{12}{(2\pi i)^4} \int \frac{z_1^{t+5} z_2^2 z_3^2 z_4^{-t-1}}{(z_1 - z_2)^3 (z_1 - z_3)^3 (z_1 - z_4)^2 (z_2 - z_3)^2 (z_2 - z_4)(z_3 - z_4)} dz \\
&= -\frac{12}{(2\pi i)^3} \int \frac{z_1^{t+5} z_2^2 z_4^{-t+1}}{(z_1 - z_2)^3 (z_1 - z_4)^5 (z_2 - z_4)^3} dz \\
&= -\frac{12}{(2\pi i)^2} \int \frac{z_1^{t+5} z_4^{-t+1}}{(z_1 - z_4)^5} \left( \frac{1}{(z_1 - z_4)^3} + 6 \frac{z_4}{(z_1 - z_4)^4} + 6 \frac{z_4^2}{(z_1 - z_4)^5} \right) dz \\
&= -12 \binom{t+5}{7} - 72 \binom{t+5}{8} - 72 \binom{t+5}{9} .
\end{aligned}$$

Adding them all up gives

$$H_4(t) = \frac{11}{11340} t^9 + \frac{11}{630} t^8 + \frac{19}{135} t^7 + \frac{2}{3} t^6 + \frac{1109}{540} t^5 + \frac{43}{10} t^4 + \frac{35117}{5670} t^3 + \frac{379}{63} t^2 + \frac{65}{18} t + 1$$

and hence

$$\text{vol } \mathcal{B}_4 = 4^3 \cdot \frac{11}{11340} = \frac{176}{2835} .$$

## 4 Larger $n$ do

As we have seen in the examples, after multiplying out the integrand of Theorem 2, many of the terms do not contribute to the integral. The following proposition will provide us with a general statement to that effect.

For a rational function  $f$  in  $n$  variables  $z_j$  we use the notation  $d_r(f)$  for the degree of  $f$  in the variables  $z_1, \dots, z_r$ .

**Proposition 3** *Suppose  $p_1, \dots, p_n$  are integers,  $q_{jk}$  are nonnegative integers ( $1 \leq j < k \leq n$ ),  $1 > \epsilon_1 > \dots > \epsilon_n > 0$ , and*

$$f(z_1, \dots, z_n) = \frac{\prod_{1 \leq j \leq n} z_j^{p_j}}{\prod_{1 \leq j < k \leq n} (z_j - z_k)^{q_{jk}}} .$$

*If  $1 \leq r \leq n$  and  $d_r(f) < -r$  then*

$$\int_{|z_1|=\epsilon_1} \dots \int_{|z_n|=\epsilon_n} f(\mathbf{z}) d\mathbf{z} = 0 .$$

*Proof.* We need only show that

$$\int_{|z_1|=\epsilon_1} \dots \int_{|z_r|=\epsilon_r} f(\mathbf{z}) d\mathbf{z} = 0 , \tag{*}$$

for then we can integrate over all  $n$  variables by first integrating over  $z_1 \dots z_r$ .

If  $r = 1$  then  $f$ , considered as a function of  $z_1$ , is analytic outside the circle  $|z_1| = \epsilon_1$  and has zero residue at infinity since its degree is less than  $-1$ , so  $(*)$  follows.

We continue by induction, so suppose  $r > 1$  and  $(*)$  is true for smaller  $r$ . Suppose that  $d_r(f) < -r$ ; we may assume  $d_{r-1}(f) \geq -(r-1)$ . Note that  $d_r(f) \geq d_{r-1}(f) + d$  where  $d$  is the degree of  $f$  in the single variable  $z_r$  (the discrepancy is the sum of the exponents  $q_{jr}$  for  $j < r$ ). Hence  $d \leq d_r(f) - d_{r-1}(f) < -1$ . We consider  $f$  as a function of  $z_r$  and apply the residue theorem to the region outside the circle  $|z_r| = \epsilon_r$ . As above  $f$  has zero residue at infinity, so we only need to consider the residues at the poles  $z_j$  for  $j < r$ . Evaluating these residues converts the integral of  $f$  into a (possibly huge) linear combination of integrals of functions of the same form as  $f$ , but in the  $n-1$  variables  $z_1, \dots, z_{r-1}, z_{r+1}, \dots, z_n$ . If  $g$  is any one of these functions then we easily calculate  $d_{r-1}(g) = d_r(f) + 1 < -(r-1)$ , and, by the induction hypothesis, the integral of  $g$  is zero.  $\square$

From this proposition we obtain the starting point for our ‘algorithm.’

**Corollary 4** For  $1 > \epsilon_1 > \dots > \epsilon_n > 0$  and  $t \geq 0$ ,  $H_n(t) =$

$$\frac{1}{(2\pi i)^n} \int_{|z_1|=\epsilon_1} \dots \int_{|z_n|=\epsilon_n} (z_1 \dots z_n)^{-t-1} \sum_{m_1+\dots+m_n=n}^* \binom{n}{m_1, \dots, m_n} \prod_{k=1}^n \left( \frac{z_k^{t+n-1}}{\prod_{j \neq k} (z_k - z_j)} \right)^{m_k} dz,$$

where  $\sum^*$  denotes that we only sum over those  $n$ -tuples of non-negative integers satisfying  $m_1 + \dots + m_n = n$  and  $m_1 + \dots + m_r > r$  if  $1 \leq r < n$ .

*Remark.* The condition on  $m_1, \dots, m_n$  can be visualized through lattice paths from  $(0, 0)$  to  $(n, n)$  using the steps  $(1, m_1), (1, m_2), \dots, (1, m_n)$ . The condition means that these paths stay strictly above the diagonal (except at the start and end).<sup>4</sup>

*Proof.* By Theorem 2,

$$\begin{aligned} H_n(t) &= \frac{1}{(2\pi i)^n} \int \dots \int (z_1 \dots z_n)^{-t-1} \left( \sum_{k=1}^n \frac{z_k^{t+n-1}}{\prod_{j \neq k} (z_k - z_j)} \right)^n dz \\ &= \frac{1}{(2\pi i)^n} \int \dots \int (z_1 \dots z_n)^{-t-1} \sum_{m_1+\dots+m_n=n} \binom{n}{m_1, \dots, m_n} \prod_{k=1}^n \left( \frac{z_k^{t+n-1}}{\prod_{j \neq k} (z_k - z_j)} \right)^{m_k} dz. \end{aligned}$$

We select a partition  $m_1 + \dots + m_n = n$  and rewrite the corresponding integrand in the language of Proposition 3:

$$p_j = (n-1)m_j + t(m_j - 1) - 1, \quad q_{jk} = m_j + m_k.$$

Now suppose  $1 \leq r < n$ . The degree of the denominator in  $z_1, \dots, z_r$  is

$$\begin{aligned} \sum_{j=1}^r \sum_{k=j+1}^n (m_j + m_k) &= (n-1) \sum_{j=1}^r m_j + r \sum_{j=r+1}^n m_j = (n-1) \sum_{j=1}^r m_j + nr - r \sum_{j=1}^r m_j \\ &= (n-1) \sum_{j=1}^r m_j + nr - r \sum_{j=1}^r (m_j - 1) - r^2. \end{aligned}$$

<sup>4</sup>The authors thank Lou Billera for pointing out this lattice-path interpretation.



We subtract this from  $\sum_{j=1}^r p_j$  to get

$$d_r(f) = (t+r) \sum_{j=1}^r (m_j - 1) - r(n-r) - r .$$

Here  $t$  is non-negative and  $r(n-r)$  is positive, so Proposition 3 implies that the integral is zero unless  $\sum_{j=1}^r (m_j - 1) > 0$ .  $\square$

Theoretically, Corollary 4 tells us what we have to do to compute  $H_n$ . Thus the integrals we computed in our calculations of  $H_3$  and  $H_4$  are exactly the non-zero integrals according to this result. For practical purposes, however, the statement is almost worthless for larger values of  $n$ . The first problem is that the number of terms in the sum equals the  $(n-1)^{\text{th}}$  Catalan number

$$\frac{1}{n} \binom{2(n-1)}{n-1} = \frac{(2n-2)!}{n!(n-1)!} ,$$

which grows exponentially with  $n$ . Another slippery point is the evaluation of each integral. As we have seen in the examples, and as can be easily seen for the general case, we can compute each integral step by step one variable at a time. However, this means at each step we convert a rational function into a sum of rational functions (of one variable less) by means of the residue theorem. Again, this means that the number of (single-variable) integrals we have to compute grows immensely as  $n$  increases. In fact, if we just ‘feed’ the statement of Corollary 4 into a computer and tell it to integrate each summand, say, starting with  $z_1$ , then  $z_2$ , and so on, the computation time explodes once one tries  $n = 7$  or  $8$ . We feel that computationally this is as involved as calculating a sufficient number of values of  $H_n$  and then interpolating this polynomial. However, complex analysis allows us some shortcuts which turn out to speed up the computation by a huge factor and which make Corollary 4 valuable, even from a computational perspective. These ‘tricks’ all showed up already in the examples and include

1. realizing when a function is analytic at the  $z_k$ -origin,
2. trying to choose the most efficient order of variables to integrate (based on estimating how many terms will be generated by the residue calculations, for each available variable), and
3. factoring the integral if some of the variables appear in a symmetric fashion.

Finally, if we are only interested in the volume of  $\mathcal{B}_n$ , we may also be

4. suppressing a particular integral if it does not contribute to the leading term of  $H_n$ .

It is worth noting that each of these computational ‘speed-ups’ decreases the total computation time substantially. By applying them to Corollary 4, we implemented a C++ program for the specific functions we have to integrate to compute  $H_n$ . We were able to verify all previously known polynomials ( $n \leq 8$ ) and to compute  $\text{vol } \mathcal{B}_9$  and  $H_9$ . The results of our calculations, including the polynomials  $H_n$  for  $n \leq 9$ , and the source code for our program are available at [www.math.binghamton.edu/dennis/Birkhoff](http://www.math.binghamton.edu/dennis/Birkhoff). The following table gives the volumes together with the respective computing time (on a 1GHz PC running under Linux). Note that computing the

full polynomial  $H_n$  takes longer, as we cannot make use of shortcut #4. In fact, the computation of  $H_9$  took about 325 days of computer time, although the elapsed time was only about two weeks since we distributed the calculations among a number of machines (8 – 40, depending on availability).

$n$	$\text{vol } \mathcal{B}_n$	time
1	1	< .01 sec
2	2	< .01 sec
3	$\frac{9}{8}$	< .01 sec
4	$\frac{176}{2835}$	< .01 sec
5	$\frac{23590375}{167382319104}$	< .01 sec
6	$\frac{9700106723}{1319281996032 \cdot 10^6}$	.18 sec
7	$\frac{77436678274508929033}{137302963682235238399868928 \cdot 10^8}$	15 sec
8	$\frac{5562533838576105333259507434329}{12589036260095477950081480942693339803308928 \cdot 10^{10}}$	54 min
9	$\frac{559498129702796022246895686372766052475496691}{2153302766311808894781211017508325066061406891577233480945238016 \cdot 10^{14}}$	317 hr

We are currently using the idle time on our departmental machines to compute the volume of the Birkhoff polytope for  $n = 10$ . We estimate the computation time to be about 10 years on one computer; our distributed calculation should be finished by the Winter of 2002. We will post the result on the Mathematics ArXiv ([front.math.ucdavis.edu](http://front.math.ucdavis.edu)).

We hope that the reader will take this immense computing time as a challenge to improve our algorithms, and work towards  $n = 11$ .

## 5 An outlook towards transportation polytopes

The Birkhoff polytopes are special cases of transportation polytopes, which are defined below. The study of this class of polytopes, which are naturally at least as fascinating as the Birkhoff polytopes, was motivated by problems in linear programming; for combinatorial properties see, for example, [10, 13]. The goal of this section is to show how our methods can be applied in this more general setting.

Fix positive real numbers  $a_1, \dots, a_m, b_1, \dots, b_n$  such that  $a_1 + \dots + a_m = b_1 + \dots + b_n$ . Let  $\mathbf{a} = (a_1, \dots, a_m)$ ,  $\mathbf{b} = (b_1, \dots, b_n)$ .

$$\mathcal{T}_{\mathbf{a}, \mathbf{b}} = \left\{ \left( \begin{array}{ccc} x_{11} & \cdots & x_{1n} \\ \vdots & & \vdots \\ x_{m1} & \cdots & x_{mn} \end{array} \right) \in \mathbb{R}^{mn} : x_{jk} \geq 0, \begin{array}{l} \sum_k x_{jk} = a_j \text{ for all } 1 \leq j \leq m, \\ \sum_j x_{jk} = b_k \text{ for all } 1 \leq k \leq n \end{array} \right\}$$

is the set of *solutions to the transportation problem with parameters  $\mathbf{a}, \mathbf{b}$* . It is a convex polytope of dimension  $(m-1)(n-1)$  in  $\mathbb{R}^{mn}$ —hence we refer to  $\mathcal{T}_{\mathbf{a}, \mathbf{b}}$  as a *transportation polytopes*. Note that  $\mathcal{B}_n = \mathcal{T}_{\mathbf{1}, \mathbf{1}}$  where  $\mathbf{1} = (1, \dots, 1) \in \mathbb{Z}^n$ . We will be exclusively interested in the case when  $a_1, \dots, a_m, b_1, \dots, b_n$  are integers. One reason for this is that the vertices of  $\mathcal{T}_{\mathbf{a}, \mathbf{b}}$  are then integral. Let

$$T(\mathbf{a}, \mathbf{b}) = T(a_1, \dots, a_m, b_1, \dots, b_n) = \#(\mathcal{T}_{\mathbf{a}, \mathbf{b}} \cap \mathbb{Z}^{mn})$$

denote the number of integer points in the transportation polytope with integral parameters  $\mathbf{a}, \mathbf{b}$ . Geometrically, each of these parameters determines the position of a hyperplane bounding the polytope  $\mathcal{T}_{\mathbf{a}, \mathbf{b}}$ . It is well known that  $T(\mathbf{a}, \mathbf{b})$  is a piecewise-defined polynomial in  $a_1, \dots, a_m, b_1, \dots, b_n$ . The regions in which  $T(\mathbf{a}, \mathbf{b})$  is a polynomial depends on the normal cones of the polytopes involved; for details, we refer to [21].

Moreover, one can derive results for the counting function  $T(\mathbf{a}, \mathbf{b})$  which are ‘higher-dimensional’ extensions of (4), by application of the main theorem in [3], which in turn is a generalization of the Ehrhart-Macdonald reciprocity theorem (2). To state this theorem, we need to define the following integer-point counting function. Suppose  $\mathcal{P}$  is a rational polytope given in the form (5); denote by  $\mathcal{P}(\mathbf{t}) = \{\mathbf{x} \in \mathbb{R}^d : \mathbf{A}\mathbf{x} = \mathbf{t}\}$  a polytope which we obtain from  $\mathcal{P} = \mathcal{P}(\mathbf{b})$  by translating (some of) its bounding hyperplanes. (Classical Ehrhart dilation is the special case  $\mathbf{t} = t\mathbf{b}$ .) Let

$$\mathcal{L}_{\mathcal{P}}(\mathbf{t}) = \#(\mathcal{P}(\mathbf{t}) \cap \mathbb{Z}^d) .$$

[3, Theorem 4] states that  $\mathcal{L}_{\mathcal{P}}$  and  $\mathcal{L}_{\mathcal{P}^\circ}$  are piecewise-defined (multivariable) quasipolynomials satisfying

$$\mathcal{L}_{\mathcal{P}}(-\mathbf{t}) = (-1)^{\dim \mathcal{P}} \mathcal{L}_{\mathcal{P}^\circ}(\mathbf{t}) .$$

As easily as (4) follows from (2), this reciprocity theorem yields a symmetry result for the transportation counting function. Let  $\mathbf{1}_d$  denote the  $d$ -dimensional vector all of whose entries are one. Then

$$T(-\mathbf{a} - n\mathbf{1}_m, -\mathbf{b} - m\mathbf{1}_n) = (-1)^{mn-1} T(\mathbf{a}, \mathbf{b}) .$$

We finally turn to the problem of writing  $T(\mathbf{a}, \mathbf{b})$  in form of an integral. As with  $\mathcal{B}_n$ , we can view  $\mathcal{T}_{\mathbf{a}, \mathbf{b}}$  as given in the form of (5) and apply the philosophy of Theorem 1 to obtain

$$T(\mathbf{a}, \mathbf{b}) = \frac{1}{(2\pi i)^{m+n}} \int \dots \int \frac{z_1^{-a_1-1} \dots z_m^{-a_m-1} w_1^{-b_1-1} \dots w_n^{-b_n-1}}{\prod_{\substack{1 \leq j \leq m \\ 1 \leq k \leq n}} (1 - z_j w_k)} dz dw .$$

Again the integral with respect to each (one-dimensional) variable is over a circle with radius  $< 1$  centered at 0, and all appearing radii are different. As in the Birkhoff case, we can separate, say, the  $\mathbf{w}$ -variables and obtain

$$T(\mathbf{a}, \mathbf{b}) = \frac{1}{(2\pi i)^{m+n}} \int \dots \int z_1^{-a_1-1} \dots z_m^{-a_m-1} \prod_{k=1}^n \int \frac{w_k^{-b_k-1}}{\prod_{j=1}^m (1 - z_j w_k)} dw_k dz .$$

And as with the Birkhoff polytope, the innermost integrals are easy to compute by means of the residue theorem. This yields the following statement which generalizes Theorem 2.

**Theorem 5** For any distinct  $0 < \epsilon_1, \dots, \epsilon_m < 1$ ,

$$T(\mathbf{a}, \mathbf{b}) = \frac{1}{(2\pi i)^m} \int_{|z_1|=\epsilon_1} \dots \int_{|z_m|=\epsilon_m} z_1^{-a_1-1} \dots z_m^{-a_m-1} \prod_{k=1}^n \sum_{j=1}^m \frac{z_j^{b_k+m-1}}{\prod_{l \neq j} (z_j - z_l)} dz.$$

*Remark.* As with Theorem 2, it can be proved from the form of the integrand that  $T(\mathbf{a}, \mathbf{b})$  is indeed a piecewise-defined polynomial in  $a_1, \dots, a_m, b_1, \dots, b_n$ .

Theoretically we could now use this theorem to produce formulas for  $T(\mathbf{a}, \mathbf{b})$  just as we did for  $H_n(t)$ . There is one major difference:  $T(\mathbf{a}, \mathbf{b})$  is only a *piecewise-defined* polynomial. In fact, we can see this from the form of the integral in Theorem 5: whether we will get a nonzero contribution at a certain step in the computation depends heavily on the relationship between  $a_1, \dots, a_m, b_1, \dots, b_n$ .

The fact that the counting function  $T(\mathbf{a}, \mathbf{b})$  is of a somewhat more delicate nature naturally has computational consequences. We feel that providing any general results on this function would go beyond the scope of this article and will hopefully be the subject of a future project. On the other hand, we adjusted our algorithm to compute values of  $T(\mathbf{a}, \mathbf{b})$  for three (fixed) pairs  $(\mathbf{a}, \mathbf{b})$ , which have been previously computed by Mount [17] and DeLoera and Sturmfels [9]. The first example is

$$T((3046, 5173, 6116, 10928), (182, 778, 3635, 9558, 11110)) = \\ 23196436596128897574829611531938753$$

The authors reported their computation took 20 minutes (Mount) or 10 minutes (DeLoera–Sturmfels) [9, 17]. We computed this number in 0.2 seconds, based on Theorem 5. A similar phenomenon happens with

$$T((338106, 574203, 678876, 1213008), (20202, 142746, 410755, 1007773, 1222717)) = \\ 316052820930116909459822049052149787748004963058022997262397$$

and

$$T((30201, 59791, 70017, 41731, 58270), (81016, 68993, 47000, 43001, 20000)) = \\ 24640538268151981086397018033422264050757251133401758112509495633028$$

which reportedly took 35 minutes/10 days with the DeLoera–Sturmfels algorithm [9], 0.3/2.9 seconds with ours. These timing comparisons ignore differences in machine speeds and implementation of the algorithms, but suggest that our methods are considerably more efficient.

**Acknowledgements.** We would like to thank the referees for carefully reading through our paper and software, and for their many helpful comments. We also thank Dean Hickerson for finding a missing factor in our original calculation of the volume of  $\mathcal{B}_9$ .

## References

- [1] Welleda Baldoni-Silva and Michèle Vergne, *Residues formulae for volumes and Ehrhart polynomials of convex polytopes*, Preprint (arXiv:math.CO/0103097), 2001.
- [2] Matthias Beck, *Counting lattice points by means of the residue theorem*, Ramanujan J. **4** (2000), no. 3, 299–310. MR 2001k:11191
- [3] ———, *Multidimensional Ehrhart reciprocity*, J. Combin. Theory Ser. A **97** (2002), no. 1, 187–194.
- [4] Louis J. Billera and A. Sarangarajan, *The combinatorics of permutation polytopes*, Formal power series and algebraic combinatorics (New Brunswick, NJ, 1994), Amer. Math. Soc., Providence, RI, 1996, pp. 1–23. MR 96m:52014
- [5] Richard A. Brualdi and Peter M. Gibson, *Convex polyhedra of doubly stochastic matrices. I. Applications of the permanent function*, J. Combinatorial Theory Ser. A **22** (1977), no. 2, 194–230. MR 55 #10486
- [6] ———, *Convex polyhedra of doubly stochastic matrices. III. Affine and combinatorial properties of  $U_n$* , J. Combinatorial Theory Ser. A **22** (1977), no. 3, 338–351. MR 55 #10488
- [7] Clara S. Chan and David P. Robbins, *On the volume of the polytope of doubly stochastic matrices*, Experiment. Math. **8** (1999), no. 3, 291–300. MR 2000k:15048
- [8] Clara S. Chan, David P. Robbins, and David S. Yuen, *On the volume of a certain polytope*, Experiment. Math. **9** (2000), no. 1, 91–99. MR 2001d:52022
- [9] Jesús DeLoera and Bernd Sturmfels, *Algebraic unimodular counting*, Preprint (arXiv:math.CO/0104286), to appear in Math. Programming B, 2002.
- [10] Persi Diaconis and Anil Gangolli, *Rectangular arrays with fixed margins*, Discrete probability and algorithms (Minneapolis, MN, 1993), Springer, New York, 1995, pp. 15–41. MR 97e:05013
- [11] Eugène Ehrhart, *Sur un problème de géométrie diophantienne linéaire. II. Systèmes diophantiens linéaires*, J. Reine Angew. Math. **227** (1967), 25–49. MR 36 #105
- [12] ———, *Sur les carrés magiques*, C. R. Acad. Sci. Paris Sér. A-B **277** (1973), A651–A654. MR 48 #10859
- [13] Victor Klee and Christoph Witzgall, *Facets and vertices of transportation polytopes*, Mathematics of the Decision Sciences, Part I (Seminar, Stanford, Calif., 1967), Amer. Math. Soc., Providence, R.I., 1968, pp. 257–282. MR 38 #4134
- [14] Donald E. Knuth, *Permutations, matrices, and generalized Young tableaux*, Pacific J. Math. **34** (1970), 709–727. MR 42 #7535
- [15] I. G. Macdonald, *Polynomials associated with finite cell-complexes*, J. London Math. Soc. (2) **4** (1971), 181–192. MR 45 #7594

- [16] Percy A. MacMahon, *Combinatory analysis*, Chelsea Publishing Co., New York, 1960. MR 25 #5003
- [17] John Mount, *Fast unimodular counting*, *Combin. Probab. Comput.* **9** (2000), no. 3, 277–285. MR 2002a:05016
- [18] Richard P. Stanley, *Linear homogeneous Diophantine equations and magic labelings of graphs*, *Duke Math. J.* **40** (1973), 607–632. MR 47 #6519
- [19] ———, *Enumerative Combinatorics*, 2nd ed., vol. I, Cambridge University Press, 1997.
- [20] M. L. Stein and P. R. Stein, *Enumeration of stochastic matrices with integer elements*, Report No. LA-4434, Los Alamos Scientific Laboratory, University of California, Los Alamos, New Mexico (1970).
- [21] Bernd Sturmfels, *On vector partition functions*, *J. Combin. Theory Ser. A* **72** (1995), no. 2, 302–309. MR 97b:52014
- [22] ———, *Equations defining toric varieties*, *Algebraic geometry—Santa Cruz 1995*, Amer. Math. Soc., Providence, RI, 1997, pp. 437–449. MR 99b:14058
- [23] Doron Zeilberger, *Proof of a conjecture of Chan, Robbins, and Yuen*, *Electron. Trans. Numer. Anal.* **9** (1999), 147–148 (electronic), *Orthogonal polynomials: numerical and symbolic algorithms* (Leganés, 1998). MR 2001b:05015

DEPARTMENT OF MATHEMATICAL SCIENCES  
 STATE UNIVERSITY OF NEW YORK  
 BINGHAMTON, NY 13902-6000  
 matthias@math.binghamton.edu  
 dennis@math.binghamton.edu